

ASSESSMENT OF VARIOUS WATER TREATMENT METHODS UTILIZED AMONG HOUSEHOLDS IN KYEGEGWA DISTRICT, UGANDA

Kiran Roberts

School of Environmental Science, University of Waterloo, Waterloo N2L 3G1, Canada
*Corresponding Author E-mail: kr Roberts@gmail.com

This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ARTICLE DETAILS

Article History:

Received 12 November 2017
Accepted 12 December 2017
Available online 1 January 2018

ABSTRACT

Safe water, coupled with adequate sanitation, and proper hygiene education (WASH) can prevent illness and death, leading to improved health, poverty reduction, and socio-economic development. The Directorate of Water Resources Management (DWRM) of MWE in Uganda, revealed that the compliance of rural safe drinking water sources to national water quality standards was 53% (E. coli being the key indicator), against a target of 95%. US \$ 147 million is lost each year due to premature deaths; approximately 23,000 Ugandans die each year from water borne diseases nearly 90% of which is directly attributed to poor water, sanitation and hygiene. The objective of this study was to assess the water treatment methods utilized among the different kinds of households in Kyegegwa district in Uganda. The research design for this study was a cross-sectional descriptive design, in which data was collected and analyzed largely by quantitative method. It used descriptive design because it offered a precise description and information of the household-based water treatments methods as they exist. Results showed that slightly more than half of the respondents used boiling as the treatment methods of choice 206 (51.9%); however, for those who used filtration majority mentioned they let it stand and settle/filter with clean cloth 80 (76.9%). For those who utilized chlorination, most of them 9 (37.5%) mentioned that they filtered 20litres and added to 2tabs of chlorine, in addition the respondents who used solar disinfection, most stated that they filtered 20litres and put in a clean container 11 (68.8%), finally majority of the respondents stated that the reason why they don't treat their water was due to the cost involved in water treatment 14 (29.8). This study concluded that the identified water treatments methods utilized among the households in Kyegegwa district were boiling, filtration, chlorination and solar disinfection.

KEYWORDS

Health, water treatment, Kyegegwa district, boiling, filtration.

1. INTRODUCTION

National Water Policy in Uganda goes thus: Separation of regulatory powers from user interests; Integrated and sustainable development, management and use of water resources with full participation of all stakeholders; Regulated use of all water other than for domestic use; Sustainable provision of clean safe water; Development and efficient use of water in agriculture; Improved collaboration and coordination among stakeholders; Equitable access and use of the Nile waters (trans-boundary water resource); Promotion of awareness of water management and development issues and capacity building; Promotion of rational, optimal and wise use of the resource for all Ugandans and all sectors; Promotion of measures for control of pollution of water resources; Promotion of the gathering and maintenance of reliable water resources information and databases; Promotion of viable management options for the resource management and provision of water supply and sanitation services[1].

The Strategies include: Establishment of an enabling environment; Development of appropriate institutional framework; Planning and prioritization of water use; Data collection and data dissemination and Capacity building. According to the water quality monitoring

work, which was carried out by the Directorate of Water Resources Management (DWRM) of MWE, the compliance of rural safe drinking water sources to national water quality standards was 53% (E. coli being the key indicator), against a target of 95%[2].

Safe water, adequate sanitation, and proper hygiene education (WSH) can prevent illness and death, leading to improved health, poverty reduction, and socio-economic development. In addition, Water is one of the key determinants to life among the element of WASH[3]. Lack of these three elements makes people more susceptible to illness and death[4].

Safe water, coupled with adequate sanitation, and proper hygiene education (WASH) can prevent illness and death, leading to improved health, poverty reduction, and socio-economic development. Household water treatment, also known as point of use water treatment (POU), such as boiling, filtration and chlorination, has been shown to be an effective means of reducing diarrhea and other diseases associated with unsafe drinking water[5].

According to a study conducted in Tanzania, 2012, which was done to determine the proportional of water treatment methods locally

available and accessible to the respondents; it was found that about 49.5% reported treating their water with any method such as boiling, strain in cloth, use of chlorine and let it stand and Settle. The Directorate of Water Resources Management (DWRM) of MWE in Uganda, revealed that the compliance of rural safe drinking water sources to national water quality standards was 53% (E. coli being the key indicator), against a target of 95%[6]. US \$ 147 million is lost each year due to premature deaths; approximately 23,000 Ugandans die each year from water borne diseases nearly 90% of which is directly attributed to poor water, sanitation and hygiene[7].

2. METHODOLOGY

2.1 Study Design

The research design for this study was a cross-sectional descriptive design, in which data was collected and analyzed largely by quantitative method. It used descriptive design because it offered a precise description and information of the household-based water treatments methods as they exist.

2.2 Study Population

The study population was 59,332 been the number of households in Kyegegwa district on accounts of studying each household.

2.3 Target Respondents

The target respondents included for the study most be aged 18 years and above with sound state of mind and should be able to speak and understand English language or the local dialects as questionnaire will be verbally translated into Rutooro/Bunyoro, Ruganda and Rukiga. The target respondent most also be living in the house unit included for study[8].

2.4 Sample Size

The sample size of the study was determined by using Slovene's formula which states that:

$$n = \frac{N}{1 + Ne^2} \quad (1)$$

where: n = Unknown sample size; N = Total population of the study.

(e) = level of significance = (0.05)²;

1 = constant;

n = 397.

2.5 Sampling Technique

The sampling technique was a multi stage cluster. The district has one County, Kyaka and nine sub-counties, namely: 1. Hapuuyo; 2. Kabweeza; 3. Kakabara; 4. Kasule; 5. Kyaka; 6. Kyegegwa town council; 7. Mpara; 8. Ruyonza and 9. Rwentuha, after clustering the sub-counties, each of the sub-counties was sampled using a simple random sampling (Table 1).

Table 1: Distribution of the sample among sub-counties of Kyegegwa District (n = 397).

District	County	Sub-county	Household (Population)	Household (Sample)
	Kyaka	Hapuuyo	9825	66
	Kyaka	Kabweeza-kyegegwa	5445	36
	Kyaka	Nakabara	10,495	70
	Kyaka	Kasule	5140	35
Kyegegwa	Kyaka	Kyaka	4398	29

Kyaka	Kyegegwa t/council	4416	30
Kyaka	Mpara	7071	47
Kyaka	Ruyonza	5043	34
Kyaka	Rwentuha	7499	50
Total		N = 59,332	n = 397

2.6 Data Collection Methods and Tools

A close ended structured researcher self-administered questionnaire for the household's. The questionnaire was prepared to cover all necessary questions about the variables under study.

A research assistant was employed for the data collection after being trained to minimized bias.

Tools will pre-test to ensure validity.

2.7 Measurements of Variables

(1) Independent variable

Attributes such as households in Kyegegwa district in Uganda were controlled by using essential questions related to each attribute.

(2) Dependent variable

Water treatment methods were measured using the following: attributes boiling, filtration, solar disinfection and chlorination.

2.8 Data Analysis and Interpretation

The data was collected and processed into tables using SPSS computer software and Microsoft excel program.

2.9 Quality Control

The research tools of this study were subjected to a pre-test, so as to assess the validity and reliability of data that would be collected for the study.

2.10 Ethical Consideration

An introductory letter from the University, addressed to the district health officer (DHO) was collected, endorsed and approved to conduct the study.

The consent to conduct the study in Kyegegwa district was sought by the district health officer (DHO).

The consent of respondent was sought, to participate in the research after citing the reasons why the research is been conducted and there right to withdraw from the study at any time of data collection process was clearly explained to respondents.

2.11 Limitations of the Study

The study encountered challenges of research assistance unwillingness to complete the questions under each questionnaire which was overcome by making sure that each questionnaire was labeled with the name of interviewer so as to be able to trace uncompleted questionnaires after data collection.

3. RESULTS

As regards to water treatment methods, it was found that slightly more than half of the respondents used boiling as the treatment methods of choice 206 (51.9%); however, for those who used filtration majority mentioned they let it stand and settle/filter with clean cloth 80 (76.9%). For those who utilized chlorination, most of them 9 (37.5%) mentioned that they filtered 20litres and added to 2tabs of chlorine, in addition the respondents who used solar disinfection, most stated that they filtered 20litres and put in a clean container 11 (68.8%), finally majority of the

respondents stated that the reason why they don't treat their water was due to the cost involved in water treatment 14 (29.8) (Table 2).

4. DISCUSSION

Water Treatment Methods Used among Households

Regarding the water treatment methods, it was found that slightly more than half of the respondents used boiling as the treatment methods of choice 206 (51.9%), followed by filtration 104 (26.2%), chlorination 24 (6.0%), solar disinfection 16 (4.0%) and those who said I don't know 47 (11.8%) respectively. However, for those who used filtration majority mentioned that they let it stand and settle/Filter with clean cloth 80 (76.9%). For those who used chlorination, most of them 9 (37.5%) mentioned that they filtered 20 liters and added to 2 tabs. For the respondents who used solar disinfection, most stated that they filtered 20 liters and put in a clean container 11 (68.8%), majority of the respondents stated that the reason why they don't treat their water was due to the cost involved in water treatment 14 (29.8). It was also seen in a slightly similar study conducted where he extracted data from the National Surveys as a retrospective study and reports on scope of household water treatment in sixty seven (67) countries worldwide indicated that the proportion of water treatment by boiling in Uganda was (39.8%) and Zambia (15.2%) whereby in Latin America chlorine is practiced by 17.1% of the households while Guinea Bissau (70.9%) and Mali (24.0%) strain drinking water through cloth[9].

Also, in a study conducted in Tanzania by Remidius Kakulu, where he determined the proportional of respondents who treat their drinking water with any method locally available and accessible to them. It was also found that about 49.5% of the respondents in the study reported treating their drinking water with any method (boiling, strain in cloth, use of chlorine/chlorination and Let it stand and Settle). Water treatment by boiling and Let it stand and Settle were frequently practiced by respondents as methods of water treatment, few of the respondents reported using water guards and strain in cloth, where boiling was the most utilized among the households which was in line to this research study[10].

Table 2: Distribution of water treatment methods used among households of Kyegegwa district.

Method	Frequency	Percent
Treatment of drinking water (method used)		
Boiling	206	51.9
Filtration	104	26.2
Chlorination	24	6.0
Solar disinfection	16	4.0
I don't treat	47	11.8
Total	397	100.0
Filtration		
Let it stand and settle/filter with clean cloth	80	76.9
All of the above	24	23.1
Total	104	100.0
Chlorination		
Filter 20 liters	7	29.2
Filter 20 liters and add to 2 tabs	9	37.5
All of the above	8	33.3
Total	24	100.0
Solar disinfection		

Filter 20 liters and Put in a clean container	11	68.8
Display in sun for 2 hrs.	5	31.2
Total	16	100.0
Reasons for not treating water		
Available	7	14.9
Cost	14	29.8
Bad taste and smelly of treatment	11	23.4
I believe water is safe from the source	10	21.3
I am used to drinking Untreated water and nothing happen to me	4	8.5
I don't know	1	2.1
Total	47	100.0

This research study stresses the need to boil and treat all water for domestic use in every households not only in the district of Kyegegwa; but should be adopted as a global measure especially in rural areas where infectious diseases tends to spread rapidly thereby causing untimely deaths and widespread poverty. However, despite this research results revealed boiling as major source of water treatment, other methods can be combined with boiling in order to kill 99.99% of bacteria that lives and reproduce in water. In addition, if water sources are consumed without any form of treatment diseases such as cholera, dysentery, typhoid, diarrhea, etc., will spread like wildfire thereby causing a population decrease due to death and economic breakdown due to loss of skills and manpower; thus water treatment should never be under looked nor taken for granted.

5. CONCLUSION

This study concluded that the identified water treatments methods utilized among the households in Kyegegwa district were boiling, filtration, chlorination and solar disinfection.

REFERENCE

- [1] Yuan, J., Lu, Y., Wang, C., Cao, X., Chen, C., Cui, H., Zhang, M., Wang, C., Li, X., Johnson, A.C., Sweetnnan, A.J., Du, D. 2020. Ecology of industrial pollution in China, *Ecosystem Health and Sustainability*, 6, (UNSP 17790101).
- [2] Perricone, V., Comi, M., Bontempo, V., Lecchi, C., Cecilian, F., Crestani, M., Ferrari, A., Savoini, G., Agazzi, A. 2020. Effects of nucleotides administration on growth performance and immune response of post-weaning piglets, *Italian Journal Of Animal Science*, 19, (1), 295-301.
- [3] Subedi, S., Roopesh, M.S. 2020. Simultaneous drying of pet food pellets and Salmonella inactivation by 395 nm light pulses in an LED reactor, *Journal of Food Engineering*, 286, (110110).
- [4] Ocheli, A., Otuya, O.B., Umayah, S.O. 2020. Appraising the risk level of physicochemical and bacteriological twin contaminants of water resources in part of the western Niger Delta region, *Environmental Monitoring and Assessment*, 192, (3245).
- [5] Geremew, A., Damtew, Y.T. 2020. Household water treatment using adequate methods in sub-Saharan countries: evidence from 2013-2016 Demographic and Health Surveys, *Journal of Water Sanitation and Hygiene For Development*, 10, (1), 66-75.
- [6] Ghadimi, F., Hajati, A., Sabzian, A. 2020. Assessment of heavy metal contamination in waters due to mineral salts company from Mighan playa/lake, Arak, Iran, *Journal of Mining and Environment*, 11, (1), 171-184.
- [7] Sudsandee, S., Fakkaew, K., Keawdounlek, V., Laor, P., Worakhunpiset, S., Apidechkul, T. 2020. Drinking water investigation of hill tribes: A case

study in northern Thailand, *International Journal of Environmental Research and Public Health*, 17, (16985).

[8] Burleson, G., Caplan, D., Mays, C., Moses, N., Navab-Daneshmand, T., Sharp, K., Maccarty, N. 2020. Computational modeling and empirical analysis of a biomass-powered drinking water pasteurization technology, *Energies*, 13, (9364).

[9] Tijani, J.O., Momoh, U.O., Salau, R.B., Bankole, M.T., Abdulkareem, A.S., Roos, W.D. 2019. Synthesis and characterization of $\text{Ag}_2\text{O}/\text{B}_2\text{O}_3/\text{TiO}_2$ ternary nanocomposites for photocatalytic mineralization of local

dyeing wastewater under artificial and natural sunlight irradiation, *Environmental Science and Pollution Research*, 26, (19SI), 19942-19967.

[10] Maju-Oyovwikowhe, G.E., Shuaib, I.I. 2019. Physiochemical characteristics and heavy metal levels of water from hand dug wells in Ikiran-Ile, Akoko Edo local government area, Edo State, Nigeria, *Journal of Applied Sciences & Environmental Management*, 23, (2), 283-290.

